

# Water refractive index in dependence on temperature and wavelength: a simple approximation

Alexey N. Bashkatov , Elina A. Genina

Optics Department Saratov State University, Saratov, Russia

## ABSTRACT

Water is the most important biological liquid. On the basis of literature data we present a simple approximation of water refractive index in dependence on temperature and wavelength in the spectral range from 200 to 1000 nm. The approximation is important for different applications in biomedical optics and optics of tissues.

**Keywords:** water refractive index, wavelength dependence, temperature dependence

## 1. INTRODUCTION

Water is the most important liquid of human organism. It is the main component of interstitial fluid, plasma of blood, intercellular fluid, etc. Knowledge of optical properties of water is very important for solution of problems of biomedical optics. In particular, wavelength dependence of the refractive index determines spectral dependence of refractive index of tissue interstitial fluid that in its turn defines the wavelength dependence of scattering properties of tissues. Furthermore, water refractive index depends on temperature.

The wavelength dependence of the water refractive index is presented in literature<sup>1,2</sup>. Wavelength dependence of water in visible and near-infrared spectral ranges has been presented as an Eq. 1<sup>1</sup>, where  $\lambda$  is wavelength, nm. In near-UV, visible and near-infrared spectral ranges wavelength dependence of water refractive index can be described by Eq. 2<sup>2</sup>. Unfortunately, these dependencies have been obtained for temperature 20 °C. At the same time, temperature of interstitial fluid is about 37 °C.

$$n(\lambda) = 1.3199 + \frac{6878}{\lambda^2} - \frac{1.132 \cdot 10^9}{\lambda^4} + \frac{1.11 \cdot 10^{14}}{\lambda^6}, \quad (1)$$

$$n(\lambda) = 1.31848 + \frac{6.662}{\lambda [nm] - 129.2}. \quad (2)$$

Taking into account the temperature dependence of water refractive index is very important for biomedical optics. Therefore, the goal of this study is obtaining the water refractive index in dependence on temperature and wavelength.

## 2. MATERIALS AND METHODS

As initial data for the construction of approximation dependencies we have used data presented in Ref. 3 (see Table 1). These data show that water refractive index decreases with increase both wavelength and temperature.

For approximation of the presented data, we have used Cauchy formula with temperature-dependent coefficients (see Eq. 3). Approximation procedure has been performed with a curve fitting system for Windows (CurveExpert, Ver. 1.34, Copyright by Daniel Hyams, 1995-1997). The Cauchy coefficients have been approximated by polynomial dependence 3-rd degree. They are presented below (see Eqs. 4-7).

---

Address all correspondence to Alexey N. Bashkatov. Tel: 8452 514693; E-mail: [bash@optics.sgu.ru](mailto:bash@optics.sgu.ru)

$$n(\lambda, t) = A(t) + \frac{B(t)}{\lambda^2} + \frac{C(t)}{\lambda^4} + \frac{D(t)}{\lambda^6}, \quad (3)$$

where  $\lambda$  is wavelength, nm; and  $A(t)$ ,  $B(t)$ ,  $C(t)$ ,  $D(t)$  are the Cauchy coefficients presented as a function of temperature;  $t$  is temperature,  $^{\circ}\text{C}$ .

Table 1. The water refractive index in dependence on temperature and wavelength<sup>3</sup>.

Temperature, $^{\circ}\text{C}$	Wavelength					
	226.5 nm	361.05 nm	404.41 nm	589 nm	632.8 nm	1013.98 nm
0	1.3945	1.34896	1.34415	1.33432	1.33306	1.32612
10	1.39422	1.3487	1.34389	1.33408	1.33282	1.32591
20	1.39336	1.34795	1.34315	1.33336	1.33211	1.32524
30	1.39208	1.34682	1.34205	1.3323	1.33105	1.32424
40	1.39046	1.3454	1.34065	1.33095	1.32972	1.32296
50	1.38854	1.34373	1.33901	1.32937	1.32814	1.32145
60	1.38636	1.34184	1.33714	1.32757	1.32636	1.31974
70	1.38395	1.33974	1.33508	1.32559	1.32438	1.31784
80	1.38132	1.33746	1.33284	1.32342	1.32223	1.31576
90	1.37849	1.33501	1.33042	1.32109	1.31991	1.31353
100	1.37547	1.33239	1.32784	1.31861	1.31744	1.31114

Relative error has been obtained from following relationship:  $\varepsilon = \frac{|n_{table} - n_{calc}|}{n_{table}}$ , where  $n_{table}$  is water refractive index presented in Table 1, and  $n_{calc}$  is water refractive index calculated with our approximation.

### 3. RESULTS AND DISCUSSION

Using the data presented in Table 1 we have approximated spectral dependence of water refractive index by the Cauchy formula for each temperature. Obtained coefficients are presented in Table 2.

Table 2. The Cauchy coefficients obtained from the calculations.

Temperature, $^{\circ}\text{C}$	A	B	$C \times 10^9$	$D \times 10^{13}$	Mean relative error $\times 10^{-3}$
0	1.310593	13417.653	-1.5969935	5.79696	1.8008839 $\times 10^{-3}$
10	1.3103755	13414.966	-1.5977596	5.8008149	1.8072102 $\times 10^{-3}$
20	1.3097143	13389.052	-1.5954159	5.7927634	1.8081556
30	1.3087499	13347.851	-1.5917125	5.7804651	1.8108928
40	1.3074721	13313.118	-1.5895477	5.7732813	1.8073181
50	1.3060093	13271.046	-1.5863232	5.76215	1.8181166
60	1.3043155	13221.979	-1.5827857	5.7498269	1.8173245
70	1.3024591	13166.14	-1.578435	5.7346675	1.8193075
80	1.3004353	13117.742	-1.575094	5.7227006	1.8321395
90	1.2982256	13059.838	-1.5709981	5.7079218	1.8310606
100	1.2959138	12995.922	-1.5660262	5.6903756	1.8432414

Using coefficients presented in Table 2 we have obtained temperature dependence of each coefficient (see Eq. 4-7).

$$A(t) = 1.3105997 - 1.4625214 \cdot 10^{-6}t - 2.3782127 \cdot 10^{-6}t^2 + 1.3505866 \cdot 10^{-8}t^3 - 4.2721445 \cdot 10^{-11}t^4 \quad (4)$$

$$B(t) = 13419.251 + 0.18056645t - 0.10699648t^2 + 0.0010813728t^3 - 4.529866 \cdot 10^{-6}t^4 \quad (5)$$

$$C(t) = -1.5972702 \cdot 10^9 - 84534.382t + 11439.948t^2 - 134.02681t^3 + 0.59353147t^4 \quad (6)$$

$$D(t) = 5.79804 \cdot 10^{13} + 3.8169 \cdot 10^9t - 4.07811 \cdot 10^8t^2 + 4.68653 \cdot 10^6t^3 - 20684.84848t^4 \quad (7)$$

### ACKNOWLEDGEMENTS

The research has been done due to financial support of grant "Leading Scientific Schools" No. 00-15-96667 of the Russian Basic Research Foundation and by Award No. REC-006 of the U.S. Civilian Research & Development Foundation for the Independent States of the Former Soviet Union (CRDF).

### REFERENCES

1. M. Kohl, M. Esseupreis, M. Cope, "The influence of glucose concentration upon the transport of light in tissue-simulating phantoms," *Phys. Med. Biol.* **40**, pp. 1267-1287, 1995.
2. P.O. Rol, *Optics for transscleral laser applications*, Ph.D. Dissertation: Swiss Federal Institute of Technology, Zurich, Switzerland. ETH N. 9665, 1991.
3. *Index of refraction of water / Handbook of Chemistry and Physics*, Eds. Lide D.R., CRC Press LLC, 2000-2001.